

## AI Abdominal Fat Measure Predicts Heart Attack and Stroke

Released: December 2, 2020

At A Glance

- An automated AI measurement of visceral fat area on abdominal CT images predicts future heart attack or stroke risk better than overall weight or BMI.
- The researchers studied 12,128 patients over 5 years.
- Visceral fat area was independently associated with future heart attack and stroke. BMI was not associated with heart attack or stroke.

OAK BROOK, Ill. — Automated deep learning analysis of abdominal CT images produces a more precise measurement of body composition and predicts major cardiovascular events, such as heart attack and stroke, better than overall weight or body mass index (BMI), according to a study presented today at the annual meeting of the Radiological Society of North America (RSNA).

[download photo](#)



Kirti Magudia, M.D., Ph.D.

“Established cardiovascular risk models rely on factors like weight and BMI that are crude surrogates of body composition,” said Kirti Magudia, M.D., Ph.D., an abdominal imaging and ultrasound fellow at the University of California San Francisco. “It’s well established that people with the same BMI can have markedly different proportions of muscle and fat. These differences are important for a variety of health outcomes.”

Unlike BMI, which is based on height and weight, a single axial CT slice of the abdomen visualizes the volume of subcutaneous fat area, visceral fat area and skeletal muscle area. However, manually measuring these individual areas is time intensive and costly.

As a radiology resident at Brigham and Women’s Hospital in Boston, Dr. Magudia was part of a multidisciplinary team of researchers, including radiologists, a data scientist and biostatistician, who developed a fully automated method using deep learning—a type of artificial intelligence (AI)—to determine body composition metrics from abdominal CT images.


“Abdominal CT scans that are routinely performed provide a more granular way of looking at body composition, but we’re not currently taking advantage of it,” Dr. Magudia said.

The study cohort was derived from the 33,182 abdominal CT outpatient exams performed on 23,136 patients at Partners Healthcare in Boston in 2012. The researchers identified 12,128 patients who were free of major cardiovascular and cancer diagnoses at the time of imaging. Mean age of the patients was 52 years, and 57% of patients were women.

The researchers selected the L3 CT slice (from the third lumbar spine vertebra) and calculated body composition areas for each patient. Patients were then divided into four quartiles based on the normalized values of subcutaneous fat area, visceral fat area and skeletal muscle area.

In this retrospective study, it was determined which of these 12,128 patients had a myocardial infarction (heart attack) or stroke within 5 years after their index abdominal CT scan. The researchers found 1,560 myocardial infarctions and 938 strokes occurred in this study group.

[download photo](#)



Michael H. Rosenthal, M.D., Ph.D.

[download photo](#)



Christopher P. Bridge, D.Phil.

Statistical analysis demonstrated that visceral fat area was independently associated with future heart attack and stroke. BMI was not associated with heart attack or stroke.

“The group of patients with the highest proportion of visceral fat area were more likely to have a heart attack, even when adjusted for known cardiovascular risk factors,” said Dr. Magudia. “The group of patients with the lowest amount of visceral fat area were protected against stroke in the years following the abdominal CT exam.”

“These results demonstrate that precise measures of body muscle and fat compartments achieved through CT outperform traditional biomarkers for predicting risk for cardiovascular outcomes,” she added.

According to Dr. Magudia, this work demonstrates that fully automated and normalized body composition analysis could now be applied to large-scale research projects.

“This work shows the promise of AI systems to add value to clinical care by extracting new information from existing imaging data,” Dr. Magudia said. “The deployment of AI systems would allow radiologists, cardiologists and primary care doctors to provide better care to patients at minimal incremental cost to the health care system.”

This paper is the recipient of an RSNA 2020 Trainee Research Prize.

Co-authors are Christopher P. Bridge, D.Phil., Camden P. Bay, Ph.D., Florian J. Fintelmann, M.D., Ana Babic, Ph.D., Katherine P. Andriole, Ph.D., Brian M. Wolpin, M.D., and Michael H. Rosenthal, M.D., Ph.D.


For more information and images, visit [RSNA.org/press20](https://www.rsna.org/press20). Press account required to view embargoed materials.

RSNA is an association of radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Illinois. ([RSNA.org](https://www.rsna.org))

Editor’s note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA media relations team at Newsroom at 1-630-590-7762.

For patient-friendly information on abdominal CT, visit [RadiologyInfo.org](https://www.radiologyinfo.org).


Video (MP4):




**Video 1.** Dr. Kirti Magudia discusses her research on automated AI measurement of visceral fat area on abdominal CT images and the prediction of future heart attack or stroke risk better than overall weight or BMI.

[Download](#)

Images (JPG, TIF):




**Figure 1.** Example of body composition analysis of an abdominal CT slice with subcutaneous fat in green, skeletal muscle in red, and visceral fat in yellow.  
[High-res \(TIF\) version](#)



**Figure 2.** Fully automated pipeline for body composition analysis using deep learning.  
[High-res \(TIF\) version](#)

# Reference Curves



*Specific to age, race and sex*

Superior to inferior percentile lines:


- 97<sup>th</sup> (red)
- 95<sup>th</sup> (orange)
- 90<sup>th</sup> (yellow)
- 75<sup>th</sup> (green)
- 50<sup>th</sup> (black)
- 25<sup>th</sup> (purple)
- 10<sup>th</sup> (pink)
- 5<sup>th</sup> (cyan)
- 3<sup>rd</sup> (blue)

Magudia et al. Radiology (in press)

**Figure 3.** Reference curves showing the distribution of body composition areas by age, race and gender.

[High-res \(TIF\) version](#)

# Heart Attack and Visceral Fat Area



Quartile 4


Quartile 3  
Quartile 2  
Quartile 1

Univariate  
 $p < 0.0001$   
Multivariate  
 $p = 0.04$

**Figure 4.** Patients in the highest quartile for normalized visceral fat area have an increased risk of subsequent heart attack.

[High-res \(TIF\) version](#)

## Stroke and Visceral Fat Area



Quartile 4

Quartile 3

Quartile 2

Quartile 1

Univariate  
 $p < 0.0001$

Multivariate  
 $p = 0.04$

**Figure 5.** Patients in the lowest quartile for normalized visceral fat area have a decreased risk of subsequent stroke.

[High-res \(TIF\) version](#)

Files:

PowerPoint presentation provided for background reporting purposes only. Do not reprint or reproduce slide content.



[Download](#)

Resources:

[Abstract\(s\) PDF](#)