

Multiple Concussions Can Disrupt Brain Connectivity in Teens

Released: November 22, 2021

At A Glance

- In a new functional MRI study, adolescents and young adults with post-concussive symptoms who suffered three to five concussions showed disruption in the brain's default mode network.
- Changes or disruptions in the default mode network have been linked to a wide number of diseases and mental disorders.
- No difference was found in the connectivity of the network in patients who had experienced only one or two concussions.

CHICAGO — Adolescents and young adults with post-concussive symptoms who suffered three to five concussions showed disruption in the default mode network, an interconnected network of brain regions involved in wakeful rest and internal thoughts. Results of study using a special MRI technique called resting-state functional MRI (fMRI) are being presented next week at the annual meeting of the Radiological Society of North America (RSNA).

[download photo](#)



Thomas Johnson, M.D., Ph.D.

"In the last 20 years, the public awareness of concussion and its long-term effects on the brain has increased and so has research into the topic," said Thomas Johnson, M.D., Ph.D., a resident physician in neurology at the University of Rochester Medical Center in Rochester, New York. "But there has been scant research using resting state fMRI at the intersection of the adolescent/young adult population and athletes with multiple concussions."

The default mode network is a functional network of brain regions that becomes active when an individual is not involved in a specific mental task but is instead daydreaming, thinking about the future or recalling memories. Brain structures included in the network are the medial prefrontal cortex, posterior cingulate cortex, inferior parietal lobule, temporal cortex and hippocampal formation. Changes or disruptions in the network have been linked to a wide number of diseases and mental disorders.

"I think of the default mode network as the idle state of the brain, similar to an idling engine," Dr. Johnson said.

The study group included 142 concussion patients (67 female, mean age 18.2 years) and age- and sex-matched controls (12 female, mean age 20.7). The patients were enrolled in a concussion clinic at the University of Rochester Medical Center for post-concussion syndrome, which includes persistent symptoms such as headaches, dizziness, fatigue and irritability.

All patients and control group participants were scanned using resting-state fMRI while doing nothing but keeping their eyes open during the imaging process. The fMRI allowed researchers to evaluate the default mode network by identifying which areas of the brain activated during imaging.

"When certain areas of the brain get active, blood flow increases in those areas," Dr. Johnson said. "Using a data analysis technique, we're able to create a map for each individual in the study that shows how activity was distributed in space and over time."

Compared to controls, the researchers found no difference in the connectivity in the default mode network in patients who had experienced one or two concussions. In patients who had three to five concussions, the maps detected a disruption in the default mode network.

"We're seeing evidence of a two-tier disease," he said. "Experiencing several concussions is a very different pathology."

Dr. Johnson said the study results give clues as to the threshold at which concussions cause effects, which in turn may help inform protocols for playing contact sports following a concussion.

"Suffering three to five concussions has the potential for long-term disruption of cognitive processes," he said. "We need to determine our tolerance for concussions. When do we say no more contact sports? We need more evidence to set some limits for people."

In the future, Dr. Johnson said he would like to conduct follow-up research on his study group by scanning the patients with resting-state fMRI in five years to determine whether disruptions in the default mode network remain.

"Trying to determine the cognitive impacts of concussion over the long-term is tricky to tease out," Dr. Johnson said. "Typically, we use standard paper tests. Using fMRI is an advanced mathematical way of looking at the networks in the brain."

In the meantime, he said resting-state fMRI may be helpful as a marker of progress and means of monitoring recovery in patients with prolonged post-concussion syndrome.

"The current standard for evaluating concussive patients is based on symptoms," Dr. Johnson said. "I envision this technique being used to help inform and guide the clinical course of patients."

Co-authors are Arun Venkataraman, Ph.D., Jianhui Zhong, Ph.D., and Edward P. Lin, M.D.

Note: Copies of RSNA 2021 news releases and electronic images will be available online at [RSNA.org/press21](https://www.rsna.org/press21).

RSNA is an association of radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Illinois. (RSNA.org)

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-791-6610.

For patient-friendly information on fMRI, visit RadiologyInfo.org.

Video (MP4):



Video 1. A radiologist reviews brain MR images.

[Download](#)



Video 2. Brain MR images displayed.

[Download](#)



Video 3. Technologist reviewing brain images.

[Download](#)



Video 4. Technologist positions a patient for brain fMRI.

[Download](#)

Images (JPG, TIF):

Multiple Concussions Can Disrupt Brain Connectivity in Teens

In a new functional MRI study, adolescents and young adults with post-concussive symptoms who suffered three to five concussions showed disruption in the brain's default mode network.



RSNA 2021
REDEFINING RADIOLOGY



Changes or disruptions in the default mode network have been linked to a wide number of diseases and mental disorders.

No difference was found in the connectivity of the network in patients who had experienced only one or two concussions.



[RSNA.org/press21](https://www.rsna.org/press21)

[High-res \(TIF\) version](#)




Figure 1. MRI images of the brain, with the default mode network (DMN) highlighted in orange/red. The DMN is active during periods of wakeful rest, when the mind is not focused on a particular task. The green and blue arrows point to areas of the network that are disrupted in adolescent patients with concussions.

[High-res \(TIF\) version](#)

Resources:

[Abstract\(s\) PDF](#)