

Ultra-high-res MRI Reveals Migraine Brain Changes

Released: November 23, 2022

At A Glance

- Researchers using ultra-high-resolution MRI have identified enlarged perivascular spaces in the brains of migraine sufferers.
- Perivascular spaces surround the blood vessels and help clear fluids from the brain
- Up to 148 million people worldwide suffer from chronic migraine.

CHICAGO — For the first time, a new study has identified enlarged perivascular spaces in the brains of migraine sufferers. Results of the study will be presented next week at the [annual meeting](#) of the Radiological Society of North America ([RSNA](#)).

"In people with chronic migraine and episodic migraine without aura, there are significant changes in the perivascular spaces of a brain region called the centrum semiovale," said study co-author Wilson Xu, an M.D. candidate at Keck School of Medicine of the University of Southern California in Los Angeles. "These changes have never been reported before."

Migraine is a common, often debilitating condition, involving a severe recurring headache. Migraines may also cause nausea, weakness and light sensitivity. According to the American Migraine Foundation, over 37 million people in the U.S. are affected by migraine, and up to 148 million people worldwide suffer from chronic migraine.

Perivascular spaces are fluid-filled spaces surrounding blood vessels in the brain. They are most commonly located in the basal ganglia and white matter of the cerebrum, and along the optic tract. Perivascular spaces are affected by several factors, including abnormalities at the blood-brain barrier and inflammation. Enlarged perivascular spaces can be a signal of underlying small vessel disease.

"Perivascular spaces are part of a fluid clearance system in the brain," Xu said. "Studying how they contribute to migraine could help us better understand the complexities of how migraines occur."

[download photo](#)



Wilson Xu

Xu and colleagues set out to determine the association between migraine and enlarged perivascular spaces.

The researchers used ultra-high-field 7T MRI to compare structural microvascular changes in different types of migraine.

"To our knowledge, this is first study using ultra-high-resolution MRI to study microvascular changes in the brain due to migraine, particularly in perivascular spaces," Xu said. "Because 7T MRI is able to create images of the brain with much higher resolution and better quality than other MRI types, it can be used to demonstrate much smaller changes that happen in brain tissue after a migraine."

Study participants included 10 with chronic migraine, 10 with episodic migraine without aura, and five age-matched healthy controls. All patients were between 25 and 60 years old. Patients with overt cognitive impairment, brain tumor, prior intracranial surgery, MRI contraindications and claustrophobia were excluded from the study.

The researchers calculated enlarged perivascular spaces in the centrum semiovale (central area of white matter) and basal ganglia areas of the brain. White matter hyperintensities—lesions that "light up" on MRI—were measured using the Fazekas scale. Cerebral microbleeds were rated with the microbleed anatomical rating scale. The researchers also collected clinical data such as disease duration and severity, symptoms at time of scan, presence of aura and side of headache.

Statistical analysis revealed that the number of enlarged perivascular spaces in the centrum semiovale was significantly higher in patients with migraine compared to healthy controls. In addition, enlarged perivascular space quantity in the centrum semiovale correlated with deep white matter hyperintensity severity in migraine patients.

"We studied chronic migraine and episodic migraine without aura and found that, for both types of migraine, perivascular spaces were bigger in the centrum semiovale," Xu said. "Although we didn't find any significant changes in the severity of white matter lesions in patients with and without migraine, these white matter lesions were significantly linked to the presence of enlarged perivascular spaces. This suggests that changes in perivascular spaces could lead to future development of more white matter lesions."

The researchers hypothesize that significant differences in the perivascular spaces in patients with migraine compared to the healthy controls might be suggestive of glymphatic disruption within the brain. The glymphatic system is a waste clearance system that utilizes perivascular channels to help eliminate soluble proteins and metabolites from the central nervous system.

However, whether such changes affect migraine development or result from migraine is unknown.

Continued study with larger case populations and longitudinal follow-up will better establish the relationship between structural changes and migraine development and type.

"The results of our study could help inspire future, larger-scale studies to continue investigating how changes in the brain's microscopic vessels and blood supply contribute to different migraine types," Xu said. "Eventually, this could help us develop new, personalized ways to diagnose and treat migraine."

Co-authors are Brendon Chou, Giuseppe Barisano, Raymond Huang, Soniya Pinto, M.D., Daniel Chang Phung, M.D., Soma Sahai-Srivastava, Alexander Lerner, M.D., and Nasim Sheikh Bahaei, M.D., FRCR.

Note: Copies of RSNA 2022 news releases and electronic images will be available online at RSNA.org/press22.

RSNA is an association of radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Illinois. (RSNA.org)

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-791-6610.

For patient-friendly information on brain MRI, visit RadiologyInfo.org.

Video (MP4):



Video 1. Technologist preparing patient for brain MRI.
[Download](#)



Video 2. Radiologist viewing brain MRI images.
[Download](#)



Video 3. Wilson Xu discusses his research on how ultra-high-res MRI reveals migraine brain changes.
[Download](#)

Ultra-high-res MRI Reveals Migraine Brain Changes

Researchers using ultra-high-resolution MRI have identified enlarged perivascular spaces in the brains of migraine sufferers.



Up to 148 million people worldwide suffer from chronic migraine.



Perivascular spaces surround the blood vessels and help clear fluids from the brain.



RSNA 2022 | Empowering Patients and Partners in Care

RSNA.org/press22

Infographic
[High-res \(TIF\) version](#)




Figure 1. (A) Cerebral microbleeds (CMB) visualized as round, dark lesions (arrow) on SWI sequence in the left temporal lobe in a migraine case with

aura. (B) Asymmetry in the appearance of the cortical vessels is more prominent on the left side (arrow) ipsilateral to the CMB.
[High-res \(TIF\) version](#)




Figure 2. (A) Punctate deep white matter hyperintensities (WMH) (arrows) on sagittal T2-weighted MRI, more prominent in the frontal lobes. (B) Migraine-free control without punctate WMH.

[High-res \(TIF\) version](#)




Figure 3. A 3D map of centrum semiovale (CSO) and basal ganglia (BG) perivascular spaces (PVS) generated from physician-reviewed automated segmentation of PVS using ITK-SNAP software.

[High-res \(TIF\) version](#)




Figure 4. (A) Enlarged centrum semiovale (CSO) perivascular spaces (PVS) (arrows) on sagittal T1-weighted MRI in a case with chronic migraine. (B) Migraine-free control without enlarged CSO PVS.

[High-res \(TIF\) version](#)



Figure 5. Technologist preparing patient for brain MRI.

[High-res \(TIF\) version](#)



Figure 6. Technologist viewing MRI images.

[High-res \(TIF\) version](#)



Figure 7. Patient undergoing brain MRI seen from control room.

[High-res \(TIF\) version](#)

Resources:

[Abstract\(s\) PDF](#)

[Your Radiologist Explains MRI Scan of the Brain](#)

[RadiologyInfo.org – Head MRI](#)

[RadiologyInfo.org – Headache](#)